

# Chapter 5

# Stochastic Processes

Econ 3070: Macroeconomics 2.0

University of Pittsburgh, 2020

# Stochastic Optimization

---

Most stochastic elements in discrete time models have straightforward analogues in continuous time

There are number of useful processes that are useful in a continuous time context because they preserve locality

The cost of introducing these is often the inclusion of higher order derivatives in value function equations

# Wiener Process

---

Bedrock process is the Wiener process / Brownian motion, characterized by  $W_0 = 0$  and

1. Independent, normal increments

$$W_s - W_t \mid W_{[0,t]} \sim \mathcal{N}(0, s - t)$$

2. Continuous with probability one

$$\lim_{h \rightarrow 0} \mathbb{P}(|W_{t+h} - W_t| > \varepsilon) = 0$$



Surprisingly few other options with just (1) as increments must be mean zero and Levy stable

- Cauchy would be another example, but this has undefined mean and variance

# Continuous Processes

---

Wiener process is the continuous limit of a random walk with increments  $\mathcal{N}(0, \Delta)$  as  $\Delta \rightarrow 0$



# Central Limit Theorem

---

It turns out that *any* standard random walk with finite mean and variance increments converges to a Wiener process

$$\frac{1}{\sqrt{N}} \sum_{i=0}^N z_i \xrightarrow{N} \text{Wiener process}$$

Note that Cauchy distribution has infinite variance so Levy Flight does not obey Central Limit Theorem

Random walks are also *recurrent*: they return to the same neighborhood infinitely often in the future. This breaks down for  $d \geq 3$  though.

# Characteristic Function

---

The characteristic function of a distribution is a spectral decomposition closely related to the Fourier transform

$$\varphi_X(t) = \mathbb{E}[\exp(itX)] = \int_{-\infty}^{\infty} \exp(itx) f(x) dx$$

This has the distinctive property that it generates the moments of the distribution

$$\mathbb{E}[X^n] = i^{-n} \left[ \frac{d^n}{dt^n} \varphi_X(t) \right]_{t=0}$$

# Spectral Convolution

---

We can also express the convolution of two variables in characteristic space. Suppose that  $X$  and  $Y$  are independent and  $Z = X + Y$  is distributed as

$$f_Z(z) = \int_{-\infty}^{\infty} f_X(x) f_Y(z - x) dx$$

In terms of characteristic functions, convolution simply becomes multiplication as in

$$\varphi_Z(t) = \varphi_X(t) \cdot \varphi_Y(t)$$

# Gaussian Basin

---

Taking a Taylor expansion, we can see that in general

$$\varphi_X(t) = \sum_{n=0}^{\infty} \frac{(it)^n}{n!} \mathbb{E}[X^n]$$

Suppose that  $\mathbb{E}[X] = 0$  and  $\mathbb{E}[X^2] = 1$ . Consider the distribution of  $Z = \frac{1}{\sqrt{n}} \sum_{i=1}^n X_i$

$$\varphi_Z(t) = \left[ \varphi_X \left( \frac{t}{\sqrt{n}} \right) \right]^n \approx \left( 1 - \frac{\frac{1}{2}t^2}{n} \right)^n \longrightarrow \exp \left( -\frac{1}{2}t^2 \right)$$

This coincides with the characteristic function of a standard normal, and so  $Z \sim \mathcal{N}(0, 1)$



# Drift Diffusion Processes

---

We can embed this into a more general class known as drift diffusion processes

$$dX_t = \mu \cdot dt + \sigma \cdot dW_t$$

And even generate a continuous time analog of an AR(1) process (the Ornstein–Uhlenbeck process)

$$dX_t = \kappa(\mu - X_t) \cdot dt + \sigma \cdot dW_t$$

# Discrete Events

---

More general drift diffusion process is represented by class of jump diffusion processes.

We can ditch the continuity assumption but preserve independent increments by bolting on a Poisson jump process.

$$dX_t = \sigma \cdot dW_t + dJ_t$$

Parameter  $\lambda$  determines jump rate. Size of jumps can even be random as long as i.i.d.

# Itô's Lemma

---

Consider the drift process  $dx = \mu dt + \sigma dW$ . For small  $\Delta$ , this evolves according to

$$x(t + \Delta) = x(t) + \Delta\mu + \sqrt{\Delta}\sigma z$$

where  $z \sim \mathcal{N}(0, 1)$

We can then approximate  $v$  by taking a Taylor expansion

$$\begin{aligned} &v(x(t + \Delta), t + \Delta) - v(x(t), t) \\ &\approx \Delta \dot{v}(x(t), t) + \mathbb{E}[(\Delta\mu + \sqrt{\Delta}\sigma z)]v_x(x(t), t) \\ &\quad + \frac{1}{2}\mathbb{E}[(\Delta\mu + \sqrt{\Delta}\sigma z)^2]v_{xx}(x(t), t) \\ &\approx \Delta \left[ \dot{v}(x(t), t) + \mu v_x(x(t), t) + \frac{\sigma^2}{2}v_{xx}(x(t), t) \right] \end{aligned}$$

# Value Function

---

The general expression for a recursive form value function is

$$\begin{aligned} v(x(t), t) &\approx \Delta u(x(t)) + \exp(-\rho\Delta) \mathbb{E} [v(x(t + \Delta), t + \Delta)] \\ \Rightarrow v(x, t) &\approx \Delta u(x) + v(x, t) \\ &+ \Delta \left[ -\rho v(x, t) + \dot{v}(x, t) + \mu v_x(x, t) + \frac{\sigma^2}{2} v_{xx}(x, t) \right] \end{aligned}$$

Eliminating  $\Delta$  and dropping explicit dependence yields the value equation

$$\rho v - \dot{v} = u(x) + \mu v_x + \frac{\sigma^2}{2} v_{xx}$$

# Distributions

---

We can apply similar techniques to distribution functions. We will denote the cumulative distribution of a variable  $x$  by  $F(x)$

Consider the case of a zero-centered O-U process with

$$dx = -\kappa(x - \mu)dt + \sigma dW$$

We can derive conditions for the evolution of the cumulative density  $F$  over  $x$ : these results are sometimes called the **Fokker-Planck** or **Kolmogorov Forward Equations**

# Discrete Limit

---

Let's approach this again using discrete approximations (note inverted signs)

$$\begin{aligned} F(x, t) &\approx \mathbb{E} \left[ F(x + \Delta\kappa(x - \mu) - \sqrt{\Delta}\sigma z, t - \Delta) \right] \\ &\approx F(x, t) - \dot{F}(x, t)\Delta + \mathbb{E}[\Delta\kappa(x - \mu) - \sqrt{\Delta}\sigma z]F_x(x, t) \\ &\quad + \frac{1}{2}\mathbb{E}[(\Delta\kappa(x - \mu) - \sqrt{\Delta}\sigma z)^2]F_{xx}(x, t) \\ &\approx F(x, t) + \Delta \left[ -\dot{F}(x, t) + \kappa(x - \mu)\sigma F_x(x, t) + \frac{\sigma^2}{2}F_{xx}(x, t) \right] \end{aligned}$$

Thus we arrive at the desired result

$$\dot{F} = \kappa(x - \mu)F_x + \frac{\sigma^2}{2}F_{xx}$$

# 0-U Limiting Distribution

---

In the limit, we will have  $\dot{F} = 0$  so that (here  $f = F_x$ )

$$\kappa(x - \mu)F_x = -\frac{\sigma^2}{2}F_{xx} \quad \Rightarrow \quad \kappa(x - \mu)f = -\frac{\sigma^2}{2}f_x$$

We can solve the prior equation using the guess  $\mathcal{N}(\mu, \gamma)$

$$f(x) = \frac{1}{\sqrt{2\pi\gamma^2}} \exp\left(-\frac{1}{2}\left(\frac{x - \mu}{\gamma}\right)^2\right)$$

Canceling common terms, this yields the following

$$\kappa = \frac{\sigma^2}{2\gamma^2} \quad \Rightarrow \quad \gamma^2 = \frac{\sigma^2}{2\kappa}$$

# Extending NCG

---

Consider the case of stochastic productivity  $z$  as an O-U process

$$f(x, k) = e^x k^\alpha$$

The derived value function is then (same FOC as before)

$$\rho v = u(e^x k^\alpha - i) + (i - \delta k)v_k - \kappa(x - \mu)v_x + \frac{\sigma^2}{2}v_{xx}$$

We can characterize the stationary distribution over  $(x, k)$  with

$$(i - \delta k)F_k = \kappa(x - \mu)F_x + \frac{\sigma^2}{2}F_{xx}$$